

“Feeding and managing ewes for improved health and efficiency”

at the farm of:

Samuel Wharry

Harphall, 42 Whitehill Road, Carnlough. BT44 0HG



Wednesday 22nd January, 2014

Researching the way forward

Feeding and managing ewes for improved efficiency

Today's farm walk aims at providing you with information and tools to inform feeding and managing decisions



Topics for discussion include:

1. Breeding strategies for a better ewe efficiency
2. Selecting for ewe prolificacy and easier-care traits
3. Diagnosis and treatment of lameness in sheep
4. Sustainable methods for liver fluke control
5. Feeding strategies for indoor lambing systems

Farm overview

Samuel Wharry, Carnlough

Farm

200 ha:

- 100 ha heather moorland (not grazed Nov-Feb)
- 25 ha rough moorland
- 60 ha unimproved grassland
- 15 ha improved grassland

Livestock

- 300 Blackface ewes + 120 crossbreds
- 100 Blackface hoggets

Sheep enterprise

- Ewes scanned end February
- All ewes lambed indoors (April-May)
- Ewes put out to hill from the end of May
- Ewes, ewe lambs and rams sold privately at home
- All other lambs fattened

Breeding programme

Samuel Wharry, Carnlough

Breeds

- Mainly Scottish Blackface ewes
- NZ Suffolk rams used for easy lambing, also Swaledale (AFBI trials)
- All replacements homebred

Ram selection

- Rams are now selected, where possible, using performance records (EBVs)
- Main criteria used in ram selection:
 - Prolificacy/maternal ability
 - Carcass quality
 - Worm resistance

Key objectives

- ◆ To breed durable ewes from within the flock with the capacity to:
 - increase numbers of lambs weaned per ewe
 - improve lambing ease
- ◆ To have a labour efficient, easier-care working system

Breeding strategies for efficient hill flocks

Breeding more efficient hill ewes

- **Ewe fertility** and **lamb growth performance** are the main constraints on profitability
- Efficiency of **crossbred ewes** shown to be equal or superior to that of purebred Blackface (BF), in particular Lleyn x BF and Swaledale x BF
- 3-way crosses: to introduce additional traits

Performance of new hill ewe types on 6 commercial flocks (1 and 2 crop ewes)


Ewe breed	Mating weight (kg)	Lambs born per 100 ewes lambled	Lambs weaned per 100 ewes lambled	Ewe efficiency (kg lamb weaned per kg ewe)
Blackface X	47	131	114	0.92
Swaledale X	48	156	136	0.96
Belclare X	53	149	120	0.91
Highlander X	50	154	131	0.97
Lleyn X	51	128	113	0.88
Texel X	53	134	110	0.82

- High level of performance (average 1.21 lambs weaned/ewe)
- Efficiencies: 15-20% higher than BF ewes (except Texel X)

Replacement breeding strategies for hill sheep flocks

Blackface ↔ Swaledale

Option 1: Criss-cross
(for 'hard' hills)




Option 2: 3-breed rotation
(for 'green' hills)

Breeding strategies for efficient lowland flocks

Breeding more efficient lowland ewes

- **Poor ewe fertility and lambing difficulties** are the main constraints on profitability
- Rotational breeding strategy: to introduce maternal traits, whilst still delivering high lamb output to market specifications



Performance of composite ewes on 6 lowland flocks (1 and 2 crop ewes)

	Ewe breed			
	Lleyn/ Belclare X	Highlander X	Romney X	Texel X
Weight at mating (kg)	59	60	60	60
Lambs weaned per 100 ewes lambled	146	167	149	135
% ewes lambled unassisted	86	89	82	88
Lamb growth rate (kg/d)	0.24	0.26	0.25	0.26
Total wt lamb weaned (kg/ewe)	48	55	52	53
Ewe efficiency (kg lamb weaned per kg ewe)	0.82	0.90	0.86	0.88

- **Highest weaning rates for Highlander X ewes**
- **Good efficiencies of 80-90%**
- **Work ongoing to assess lifetime performance**

Selecting ewes for maternal traits

Hillsborough Management Recording Scheme

◆ Objectives:

- Identify ewes in commercial flocks suited to easier-care systems
- Breed replacement sheep that will require less intervention at lambing in future easier-care systems



◆ Recording & analysis:

- Step 1: Simple recording of key traits for prolificacy and easier-care traits (provision of summary report)
- Step 2: Simple recording of lamb live weights (provision of performance index of ewes)

Ewes and replacements ranked on performance on a scale 0-100

Step 1: Lambing book

Key traits for 'easier management':

Lambing ease →

Mothering ability →

Lamb viability →

Ewe no:	52	Date of lambing:	21/03/05
Ewe breed:	B	Sire breed (ID)	LL
Ewe details			
Age at lambing:	1-yr	2-yr	3yr+
Lambing difficulty score:	No help	Little help	Manual delivery: OK Difficult
If helped Why?	Management	Oversized	Malpresented
Mothering ability:	Follows whatever	Stands well back	Leaves lambs
Lamb details			
Lamb tag no:	22	23	24
Lamb sex:	M F	M F	M F
Fostered to:	Ewe no	Ewe no	Ewe no
Lamb viability:	Up & suck	Slow suck	Help suck
Date of mortality:			
General ewe problems			
Teat problems:		Yes	
Insufficient colostrum:		Yes	
Prolapse:		Yes	

Step 2: Lamb live weight book

Key traits for productivity:
number of lambs reared
per ewe and lamb weight

From lambing book

Farm name		2010
Ewe Details		% of total
Total number of ewes	84	
Average number of lambs per ewe	1.5	145.2
No. of ewes with 1 lamb	43	51.2
No. of ewes with 2 lambs	34	40.5
No. of ewes with 3 lambs	7	8.3
No. of ewes lambled unaided	55	65.5
No. of ewes that needed some help	18	21.4
No. of ewes that needed manual help	9	10.7
Unavailable data	2	2.4
No. of ewes who follows lamb whatever	73	86.9
No. of ewes who stands well back	2	2.4
Unavailable data	9	10.7
Lamb Details		% of total
Total number of lambs	122	
Number of lambs born dead	1	0.8
Number of lambs born alive	121	99.2
No. of lambs up to suck	116	95.9
No. of lambs slow to suck	0	0.0
No. of lambs needing help to suck	5	4.1

From lamb live weight book

Farm name		2011					
Ewe	Sire	Age	No lambs	LDS	MA	WWT	INDEX
281	BL	3	2	1	1	77	100
304	T(P)	2	2	1	1	92	98
211	S(M)	3	2	1	1	92	87
310	T(F)	2	3	1	1	113	86
272	T(G)	3	3	1	1	119	85
109	T(P)	3	2	1	1	77	84
286	S(B)	3	2	1	1	89	84
207	S(M)	3	2	1	1	85	83
153	S(B)	3	2	1	1	88	83
287	S(B)	3	2	1	1	85	82
200	T(F)	3	2	1	1	92	81
302	T(P)	2	2	2	1	92	81
280	T(F)	3	2	1	1	91	80
140	T(F)	3	2	1	1	90	80
251	S(T)	3	2	1	1	84	79
177	T(V)	3	2	1	1	78	77
94	S(W)	3	2	2	1	78	74
205	S(M)	3	2	2	1	98	73
265	T(G)	3	3	2	1	127	72
326	S(M)	2	2	2	1	81	70
...

Case study

Average number of ewes in the flock: 85 (mostly Blackface)

	2007	2008	2010	2013	Trend
Easier management traits					
% ewes lambed unaided	65	55	67	80	+
% ewes who follows lamb	77	72	97	97	+
% lambs up to suck	93	95	96	94	+
Productivity traits					
% ewes with > 1 lamb	42	41	49	47	+
% lambs born alive	97	98	99	98	+

Diagnosis and treatment of lameness in sheep

Do you know the cause ?

- Lameness can cause long-term pain and increase production costs (due to reduced feed intake) and treatment costs
- Knowing the cause of lame sheep is the first step towards its treatment, control and prevention
- Main issues identified in NI sheep flocks surveyed:

Shelly hoof



Scald



Footrot




Toe granulomas ('strawberry')



Towards a better control

Booklet available to:

- Better diagnose the cause
- Identify appropriate treatment options
- Know how to prevent the conditions
- Follow best practice for foot bathing, foot trimming



Diagnosis and treatment of lameness in sheep


Treatment and prevention

Key points

- Always separate lame sheep and treat last
- Always record or mark treated animals
- Seek veterinary advice if necessary
- In most cases, routine trimming of all feet is unnecessary
- When foot trimming, clean and disinfect foot shears and treatment area, and dispose of any hoof trimmings
- After treatments, animals should stand on a hard, clean and dry surface to maximise efficacy

The 'Stamp out lameness' Campaign


1. Cull badly or repeatedly infected animals
2. Quarantine incoming animals
3. Correct diagnosis and prompt treatment
4. Avoid spreading infection at handling and gathering
5. Adopt a footrot vaccination program



Treatment of Liver Fluke in Sheep

Key Issues

- ◆ Choose the right product
- ◆ Re-infection and re-treatment
– no residual effect
- ◆ Use of adulticides
- ◆ Avoid unnecessary use of combinations
- ◆ Correct dose rate, drenching /
application technique



Indoor lambing system

Feeding strategy at Samuel Wharry's Farm

- All concentrate diet, no silage
- Ewes scanned in February
- Twin-bearing ewes housed post scanning
- Lambing starts 4th of April
- Ewes have access to straw *ad lib*

Feed rates (kg/day)		
Weeks to lambing	Singles	Twins
8 to 4	0.0	1.0
4 to 1	0.25	1.0
1 to lambing	0.7	1.0

- Blend of Maize, barley, soya hulls, distillers dark grain, citrus pulp, soya meal, molasses, minerals and vitamins

Indoor lambing system

Silage and concentrate diets

- ◆ Produce high quality silage to reduce concentrate input
- ◆ Match concentrate requirements with forage quality to ensure
 - Viable lambs
 - Udder development
 - Adequate colostrum
 - Maternal bonding

Silage Quality		
	Good	Poor
Dry-matter %	28.4	13.3
ME (MJ/kg DM)	11.2	8.4
CP (% DM)	12.1	8.2
D-value (% DM)	67.3	56.4
Concentrate feed over 6 weeks (kg)	12	28

- ◆ Feed rate driven by forage quality
- ◆ Must be cost effective

Indoor lambing system

Concentrate supplementation

Key issues:

- ◆ Assess nutritional status of ewes and establish litter size
- ◆ Know the feeding value of your silage
- ◆ Consider the pattern and frequency of meal feeding
- ◆ Concentrate composition is important



Energy: - Target 11-12 MJ/kg
- Cereals: feed whole with hay or processed with silage
- Beet pulps/soya hulls

Protein: - Target 17-21% CP & 45-55 g/kg DUP

Vit/Min: - Selenium 0.2-0.4 mg/kg
- Vitamin E 100-150 IU/kg

Benchmarking Farm Performance

Samuel Wharry, Carnlough

Physical performance

	2011/12	2012/13		2012/13 Average
Number of ewes	365	347		311
Lambs sold/ewe	1.08	1.19		1.28
Concentrates fed (kg/ewe)	105	146		74
Av. carcass weight (kg)	21	21		20
Kg carcass/ha	29	34		84

Benchmarking Farm Performance

Samuel Wharry, Carnlough

Financial performance (£/ewe)

	2011/12	2012/13		2012/13 Average
Lamb sales	82	96		75
Replacement cost	-12	-16		-14
Total output	62	79		68
Total variable costs	52	55		47
Gross margin	10	23		20

Researching the way forward

Other current sheep research projects

- Interrelationships between trace element status, gastrointestinal parasite infection and growth performance of lambs
- Development of sustainable livestock systems to promote biodiversity within hill areas (by identifying breeding and grazing strategies)
- Factors affecting eartag retention in sheep
- Effects of breed and forage type on methane emissions from sheep
- Meat quality of entire male versus castrate lambs finished on forage-based diets



This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.